arXiv:cond-mat/0612653v1 [cond-mat.mes-hall] 27 Dec 2006

Square-gradient mechanism of surface scattering in quasi-1D rough waveguides

F. M. Izrailev
Instituto de Física, Universidad Autónoma de Puebla,
Apartado Postal J-48, Puebla, Pue., 72570, México

N. M. Makarov*
A. Ya. Usikov Institute for Radiophysics and Electronics,
Ukrainian Academy of Science, 12 Proskura Street, 61085 Kharkov, Ukraine

M. Rendón
Facultad de Ciencias de la Electrónica, Universidad Autónoma de Puebla, Puebla, Pue., 72570, México

(Dated: February 6, 2008)

We develop a perturbative approach that allows one to study the surface scattering in quasi-1D waveguides with rough boundaries. Our approach is based on the construction of an effective "bulk" potential of a very complicated structure. The detailed analysis of this potential reveals that apart from the well known terms considered in the previous studies, one should keep specific terms that depend on the square of the derivative of the boundaries. As was found, in spite of an apparent smallness of these *square-gradient scattering* (SGS) terms, there is a physically important region of parameters, in which they can not be neglected. Our approach also demonstrates that the contribution of the SGS-mechanism of scattering strongly depends on the type of the rough boundaries (uncorrelated, symmetric or antisymmetric).

PACS numbers: 73.21.Hb, 42.25.Dd

I. INTRODUCTION

Wave scattering in surface-corrugated guiding systems remains a quite interesting problem, both from the theoretical viewpoint and due to many applications ranging from the sound and electromagnetic propagation, to the conduction of electrons in mesoscopic wires. In spite of many important results obtained to date (see, e.g., Refs. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]), there is a number of open questions to be resolved. One of such problems is to understand the interplay between different mechanisms of the scattering that are due to principally different terms in an effective scattering potential constructed with the use of various perturbative approaches.

In the recent papers [17] it was shown that apart from the well known terms in the potentials appearing in the first order perturbation theory, one should keep some specific terms that were neglected due to their seeming smallness. It was demonstrated that these terms, although formally belonging to the second order terms, can play a decisive role in the scattering in some region of parameters. Since these terms are proportional to the square of the derivative of surface profiles, they can be associated with the *square-gradient scattering* (SGS) mechanism.

In contrast with Ref.[17] where a particular case of one rough surface was considered, below we develop a general approach by concentrating our attention on quasi-one dimensional waveguides with two corrugated surfaces that are either statistically uncorrelated or correlated. Our main interest is in the expression for the *attenuation length* L_n (also known as the scattering length, or the total mean free path), corresponding to specific n-th propagating mode. We assume that the total number N of these modes (or conducting channels) is large, $N \gg 1$, and that the lower and upper surface profiles (boundaries) of such quasi-1D waveguides are described by two random functions $\xi_\downarrow(x)$ and $\xi_\uparrow(x)$, respectively.

In what follows, we consider three configurations of a particular interest:

1. The waveguide with the *uncorrelated* boundaries, $\xi_\downarrow(x)$ and $\xi_\uparrow(x)$.
2. The waveguide with the *antisymmetric* boundaries,
$$\xi_\downarrow(x) = \xi_\uparrow(x) = \xi(x). \qquad (1.1)$$
3. The waveguide with the *symmetric* boundaries,
$$-\xi_\downarrow(x) = \xi_\uparrow(x) = \xi(x). \qquad (1.2)$$

In our approach we assume that due to the multiple scattering of a travelling wave from the rough boundaries, the longitudinal wave number of an n-th propagating mode can be written as $k_n + \delta k_n$, where k_n is its unperturbed value,

$$\delta k_n = \gamma_n + i(2L_n)^{-1}. \qquad (1.3)$$

---

[*] On sabbatical leave from Instituto de Ciencias, Universidad Autónoma de Puebla, Priv. 17 Norte No. 3417, Col. San Miguel Hueyotlipan, Puebla, Pue., 72050, México

The real part γ_n is responsible for a roughness-induced correction to the phase velocity of a given mode. As is known, the shift γ_n does not change the transport properties of a disordered system. Therefore, our further analysis shall be focused on the attenuation length L_n only.

In order to properly take into account the surface irregularities we use the approach developed in Ref. [17]. Specifically, by means of the canonical transformation we formally reduce the problem of the *surface* scattering to the *bulk* one described by the Hamiltonian $\widehat{H}$,

$$\widehat{H} = \widehat{H}^{(0)} + \widehat{U} \tag{1.4}$$

with the effective potential $\widehat{U}$. It should be stressed that such a transformation does not mean an equivalence between the surface and bulk scattering. The very point is that the obtained "bulk" Hamiltonian $\widehat{H}$ is of very specific form that does not allow to use standard random matrix approaches based on a complete randomness in effective potentials.

The paper is organized as follows. In Sec. II we start with the general expressions describing the waveguide with any form of rough boundaries. In Sec. III we write the Dyson-type equation and derive approximate expressions for the surface scattering potential in the form, most useful for our perturbative approach. In Sec. IV we obtain the fluctuating part of the potential for three configurations of a waveguide. In Sec. V we present a brief description of the formalism for obtaining the average Green's function developed in Ref. [17]. Here we also specify the conditions of applicability of our method. In Sec. VI we perform a detailed analysis of the attenuation length for three specific configurations of the waveguide. In Sec. VII we report the data that illustrates our analytical results and make clear the role of various terms in the expression for the scattering potential. In this Section we also discuss the so-called "repulsion" phenomenon discovered in Ref. 18, in view of its relation to the square-gradient scattering mechanism. A brief comparison of the three configurations, as well as our concluding remarks, are presented in Sec. VIII.

II. THE WAVEGUIDE WITH GENERAL ROUGH BOUNDARIES

In this section we derive the general expressions describing quasi-1D waveguides with any type of the boundaries. Then, we present them in the form, most suitable for our analysis of three particular waveguide configurations.

Specifically, we consider the standard model of an open waveguide of the average width d, stretched along the x axis. The lower and upper surfaces of the waveguide are assumed to be described, respectively, by the profiles $z = \sigma\xi_{\downarrow}(x)$, and $z = d+\sigma\xi_{\uparrow}(x)$. Here σ is the root-mean-square roughness height that is assumed to be identical for both boundaries. In other words, the waveguide occupies the region

$$-\infty \le x \le \infty, \qquad \sigma\xi_{\downarrow}(x) \le z \le d + \sigma\xi_{\uparrow}(x) \tag{2.1}$$

of the (x, z)-plane. The fluctuating width $w(x)$ is defined by

$$w(x) = d + \sigma[\xi_{\uparrow}(x) - \xi_{\downarrow}(x)], \qquad \langle w(x)\rangle = d\,, \tag{2.2}$$

where the random functions $\xi_i(x)$ (with $i =\downarrow, \uparrow$) describe the roughness of the corresponding boundary.

In what follows we consider the model for which both of the boundaries are statistically identical. They are assumed to be described by a statistically homogeneous and isotropic Gaussian random processes with the zero mean and unit variance,

$$\langle \xi_i(x)\rangle = 0, \quad \langle {\xi_i}^2(x)\rangle = 1, \tag{2.3a}$$

$$\langle \xi_i(x)\xi_i(x')\rangle = \mathcal{W}(x - x'). \tag{2.3b}$$

Here the angular brackets stand for the statistical averaging over different realizations of the surface profiles $\xi_i(x)$. We also assume that the binary correlator $\mathcal{W}(x)$ decreases on the scale R_c, with the normalization, $\mathcal{W}(0) = 1$.

For three configurations we analyze below, the cross-correlations are as follows. In the first case of the uncorrelated boundaries we obviously have,

$$\langle \xi_{\downarrow}(x)\xi_{\uparrow}(x')\rangle = 0. \tag{2.4}$$

The waveguide with the antisymmetric boundaries implies:

$$\langle \xi_{\downarrow}(x)\xi_{\uparrow}(x')\rangle = \mathcal{W}(x - x'). \tag{2.5}$$

As for the symmetric boundaries case, the opposite boundaries hold the correlation equality:

$$\langle \xi_{\downarrow}(x)\xi_{\uparrow}(x')\rangle = -\mathcal{W}(x - x'). \tag{2.6}$$

The *roughness-height power* (RHP) spectrum $W(k_x)$ is defined by:

$$W(k_x) = \int_{-\infty}^{\infty} dx \exp(-ik_x x)\, \mathcal{W}(x). \tag{2.7}$$

Since $\mathcal{W}(x)$ is an even function of x, its Fourier transform (2.7) is an even, real and non-negative function of k_x. The RHP spectrum has the maximum at $k_x = 0$ with $W(0) \sim R_c$, and decreases on the scale R_c^{-1}.

In order to analyze the surface scattering problem we shall employ the method of retarded Green's function $\mathcal{G}(x, x'; z, z')$. Specifically, we start with the Dirichlet boundary-value problem

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial z^2} + k^2\right)\mathcal{G}(x, x'; z, z')$$

$$= \delta(x-x')\delta(z-z'), \qquad (2.8a)$$

$$\begin{aligned} &\mathcal{G}(x,x';z=\sigma\xi_\downarrow(x),z')=0\\ &\mathcal{G}(x,x';z=d+\sigma\xi_\uparrow(x),z')=0. \end{aligned} \qquad (2.8b)$$

Here $\delta(x)$ and $\delta(z)$ are the Dirac delta functions. The wave number k is equal to ω/c for an electromagnetic wave of the frequency ω and TE polarization, propagating through a waveguide with perfectly conducting walls. For an electron quantum wire, k is the Fermi wave number within the isotropic Fermi-liquid model.

A. Effective surface scattering potential, $\widehat{U}$

The equation (2.8a) does not contain any scattering potential, therefore, the perturbation is hidden in the boundary conditions (2.8b). In this case it is convenient to perform the canonical transformation to new coordinates,

$$x_{new} = x_{old}, \qquad (2.9a)$$

$$\begin{aligned} z_{new} &= \frac{d\,[z_{old}-\sigma\xi_\downarrow(x_{old})]}{w(x_{old})}\\ &= \frac{d\,[z_{old}-\sigma\xi_\downarrow(x_{old})]}{d+\sigma[\xi_\uparrow(x_{old})-\xi_\downarrow(x_{old})]}, \end{aligned} \qquad (2.9b)$$

in which both boundaries are flat. In accordance with Eqs. (2.9), the transformations for the first and second derivatives with respect to the old coordinates are given by (below for convenience we drop the subscript *new* for x and z):

$$\begin{aligned} \frac{\partial}{\partial x_{old}} &= \frac{\partial x}{\partial x_{old}}\frac{\partial}{\partial x}+\frac{\partial z}{\partial x_{old}}\frac{\partial}{\partial z}\\ &= \frac{\partial}{\partial x}-\left[\frac{\sigma\xi'_\uparrow(x)}{w(x)}z+\frac{\sigma\xi'_\downarrow(x)}{w(x)}(d-z)\right]\frac{\partial}{\partial z}, \end{aligned} \qquad (2.10a)$$

$$\begin{aligned} \frac{\partial}{\partial z_{old}} &= \frac{\partial x}{\partial z_{old}}\frac{\partial}{\partial x}+\frac{\partial z}{\partial z_{old}}\frac{\partial}{\partial z}\\ &= \frac{d}{w(x)}\frac{\partial}{\partial z}, \end{aligned} \qquad (2.10b)$$

$$\begin{aligned} \frac{\partial^2}{\partial x_{old}^2} &= \left\{\frac{\partial}{\partial x}-\left[\frac{\sigma\xi'_\uparrow(x)}{w(x)}z+\frac{\sigma\xi'_\downarrow(x)}{w(x)}(d-z)\right]\frac{\partial}{\partial z}\right\}\\ &\times \left\{\frac{\partial}{\partial x}-\left[\frac{\sigma\xi'_\uparrow(x)}{w(x)}z+\frac{\sigma\xi'_\downarrow(x)}{w(x)}(d-z)\right]\frac{\partial}{\partial z}\right\} \end{aligned} \qquad (2.11a)$$

$$\frac{\partial^2}{\partial z_{old}^2} = \frac{d^2}{w^2(x)}\frac{\partial^2}{\partial z^2}. \qquad (2.11b)$$

Correspondingly, we introduce the canonically conjugate Green's function,

$$\mathcal{G}_{new}(x,x';z,z')=\frac{\sqrt{w(x)w(x')}}{d}\,\mathcal{G}_{old}(x,x';z,z'). \qquad (2.12)$$

Here the prefactor before $\mathcal{G}_{old}$ is due to the Jacobian of the transformation to new variables. As a result, we arrive at the equivalent boundary-value problem governed by the equation,

$$\begin{aligned} &\left(\frac{\partial^2}{\partial x^2}+\frac{\partial^2}{\partial z^2}+k^2\right)\mathcal{G}(x,x';z,z')\\ &-\widehat{U}(x,z)\,\mathcal{G}(x,x';z,z')=\delta(x-x')\delta(z-z'), \end{aligned} \qquad (2.13a)$$

$$\begin{aligned} &\mathcal{G}(x,x';z=0,z')=0,\\ &\mathcal{G}(x,x';z=d,z')=0. \end{aligned} \qquad (2.13b)$$

Here we also omitted the subscript *new* for $\mathcal{G}_{new}(x,x';z,z')$. As a result of the transformation, we have formally reduced the problem of surface scattering to that of the bulk one specified by the effective potential $\widehat{U}(x,z)$,

$$\begin{aligned} \widehat{U}(x,z) &= \left[1-\frac{d^2}{w^2(x)}\right]\frac{\partial^2}{\partial z^2}\\ &+\frac{\sigma}{w(x)}\Bigg\{\left[\xi''_\uparrow(x)+2\xi'_\uparrow(x)\frac{\partial}{\partial x}\right]\left[\frac{1}{2}+z\frac{\partial}{\partial z}\right]\\ &-\left[\xi''_\downarrow(x)+2\xi'_\downarrow(x)\frac{\partial}{\partial x}\right]\left[\frac{1}{2}+(z-d)\frac{\partial}{\partial z}\right]\Bigg\}\\ &-\frac{\sigma^2}{w^2(x)}\Bigg\{{\xi'_\uparrow}^2(x)\left[\frac{3}{4}+3z\frac{\partial}{\partial z}+z^2\frac{\partial^2}{\partial z^2}\right]\\ &+{\xi'_\downarrow}^2(x)\left[\frac{3}{4}+3(z-d)\frac{\partial}{\partial z}+(z-d)^2\frac{\partial^2}{\partial z^2}\right]\\ &-\xi'_\downarrow(x)\,\xi'_\uparrow(x)\left[\frac{3}{2}+3(2z-d)\frac{\partial}{\partial z}+2z(z-d)\frac{\partial^2}{\partial z^2}\right]\Bigg\}. \end{aligned} \qquad (2.14)$$

Note that the prime over the function $\xi_i(x)$ denotes a derivative with respect to x. It is important to stress that Eqs. (2.13) and (2.14) are *exact* ones, and valid for any form of the surface profiles $\xi_\downarrow(x)$ and $\xi_\uparrow(x)$.

As is expected, the general expression (2.14) reduces to that obtained previously in Ref.[17] for a particular case of one rough boundary, if one substitutes $\xi_\downarrow(x)=0$ and $\xi_\uparrow(x)=\xi(x)$.

III. DYSON EQUATION

Here we start with the non-local Dyson equation in k_x-representation,

$$g(k_x,k'_x)=2\pi\delta(k_x-k'_x)\,g_0(k_x)$$

$$+\,g_0(k_x)\int_{-\infty}^{\infty}\frac{dq_x}{2\pi}\,\Xi(k_x,q_x)g(q_x,k'_x) \qquad (3.1)$$

with $g_0(k_x)=k_z\cot(k_z d)$ being the unperturbed pole-factor. In this representation the effective surface scattering potential $\Xi(k_x,k'_x)$ is given by the expression,

$$\begin{aligned}\Xi(k_x,k'_x)=&\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\,\exp\left(-ik_x x_1\right)\frac{\sin(k_z z_1)}{k_z}\\ &\times\widehat{U}(x_1,z_1)\exp\left(ik'_x x_1\right)\frac{\sin(k'_z z_1)}{k'_z}\,. \qquad (3.2)\end{aligned}$$

Here k_x and k_z are the lengthwise and transverse wave numbers,

$$k_z=k_z(k_x)=\sqrt{k^2-k_x^2},\qquad k'_z=k_z(k'_x). \qquad (3.3)$$

Their unperturbed eigenvalues are k_n,

$$k_n=\sqrt{k^2-(\pi n/d)^2},\qquad n=1,2,3,\ldots,N_d. \qquad (3.4)$$

and $\pi n/d$, respectively. The total number N_d of propagating waveguide modes (conducting channels that have real values of k_n) is determined by the integer part $[\ldots]$ of the ratio kd/π,

$$N_d=[kd/\pi]. \qquad (3.5)$$

After the substitution of the expression (2.14) for $\widehat{U}$ in Eq. (3.2), we realize that $\Xi(k_x,k'_x)$ consists of three groups of terms of a different nature. The first one has the factor $[1-d^2/w^2(x_1)]$, the second and third groups have the factors $\sigma/w(x_1)$ and $\sigma^2/w^2(x_1)$, respectively. One needs to note that while itself the kernel $\Xi(k_x,k'_x)$ is Hermitian, the latter two parts are non-Hermitian ones. In order to present each part of $\Xi(k_x,k'_x)$ in an Hermitian form, we perform the integration by parts for the term containing $\xi''_{\downarrow}(x_1)$ and $\xi''_{\uparrow}(x_1)$ in the second group. As a result, we arrive at the final *exact* form for the perturbation potential,

$$\begin{aligned}\Xi(k_x,k'_x)=&\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\,\exp\left(-ik_x x_1\right)\frac{\sin(k_z z_1)}{k_z}\\ &\times\Bigg(\left[1-\frac{d^2}{w^2(x_1)}\right]\frac{\partial^2}{\partial z_1^2}+\frac{\sigma}{w(x_1)}\left[i(k_x+k'_x)\right]\\ &\times\left\{\xi'_{\uparrow}(x_1)\left[\frac{1}{2}+z_1\frac{\partial}{\partial z_1}\right]-\xi'_{\downarrow}(x_1)\left[\frac{1}{2}+(z_1-d)\frac{\partial}{\partial z_1}\right]\right\}\\ &-\frac{\sigma^2}{w^2(x_1)}\Bigg\{{\xi'_{\uparrow}}^2(x_1)\left[\frac{1}{4}+2z_1\frac{\partial}{\partial z_1}+z_1^2\frac{\partial^2}{\partial {z_1}^2}\right]\\ &+{\xi'_{\downarrow}}^2(x_1)\left[\frac{1}{4}+2(z_1-d)\frac{\partial}{\partial z_1}+(z_1-d)^2\frac{\partial^2}{\partial {z_1}^2}\right]\\ &-\xi'_{\downarrow}(x_1)\,\xi'_{\uparrow}(x_1)\Bigg[\frac{1}{2}+2(2z_1-d)\frac{\partial}{\partial z_1}\\ &+2z_1(z_1-d)\frac{\partial^2}{\partial {z_1}^2}\Bigg]\Bigg\}\Bigg)\exp\left(ik'_x x_1\right)\frac{\sin(k'_z z_1)}{k'_z} \qquad (3.6)\end{aligned}$$

This equation has a peculiar structure, very useful for a further analysis. The kernel written in this form consists also of three groups of terms, however, now they can be associated with different scattering mechanisms.

Since we are interested in the averaged Green's function, we have to calculate the binary correlator of $\Xi(k_x,k'_x)$. Therefore, in order to avoid very cumbersome calculations, it is reasonable to make a simplification of $\Xi(k_x,k'_x)$ that does not destroy the Hermitian structure of each group of terms. Taking also into account that our main interest is in the case of small surface corrugations $\sigma\ll d$, we can do the following: expand the factor $1-d^2/w^2(x_1)\approx 2\sigma[\xi_{\uparrow}(x_1)-\xi_{\downarrow}(x_1)]/d$ in the first term of Eq. (3.6) and to put $w(x_1)\approx d$ in all the others. In such a way, we derive the following *approximate* expression for the surface scattering potential,

$$\begin{aligned}\Xi(k_x,k'_x)\approx&\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\,\exp\left(-ik_x x_1\right)\frac{\sin(k_z z_1)}{k_z}\\ &\times\Bigg(\frac{2\sigma}{d}\left[\xi_{\uparrow}(x_1)-\xi_{\downarrow}(x_1)\right]\frac{\partial^2}{\partial z_1^2}+\frac{\sigma}{d}\left[i(k_x+k'_x)\right]\\ &\times\left\{\xi'_{\uparrow}(x_1)\left[\frac{1}{2}+z_1\frac{\partial}{\partial z_1}\right]-\xi'_{\downarrow}(x_1)\left[\frac{1}{2}+(z_1-d)\frac{\partial}{\partial z_1}\right]\right\}\\ &-\frac{\sigma^2}{d^2}\Bigg\{{\xi'_{\uparrow}}^2(x_1)\left[\frac{1}{4}+2z_1\frac{\partial}{\partial z_1}+z_1^2\frac{\partial^2}{\partial {z_1}^2}\right]\\ &+{\xi'_{\downarrow}}^2(x_1)\left[\frac{1}{4}+2(z_1-d)\frac{\partial}{\partial z_1}+(z_1-d)^2\frac{\partial^2}{\partial {z_1}^2}\right]\\ &-\xi'_{\downarrow}(x_1)\,\xi'_{\uparrow}(x_1)\Bigg[\frac{1}{2}+2(2z_1-d)\frac{\partial}{\partial z_1}\\ &+2z_1(z_1-d)\frac{\partial^2}{\partial {z_1}^2}\Bigg]\Bigg\}\Bigg)\exp\left(ik'_x x_1\right)\frac{\sin(k'_z z_1)}{k'_z} \qquad (3.7)\end{aligned}$$

The expression (3.7) contains three types of scattering terms of a different physical meaning: the terms depending on the roughness height $\sigma\xi_i(x_1)$, derivative of the roughness height $\sigma\xi'_i(x_1)$, and square of the derivative of the roughness height $\sigma^2{\xi'_i}^2(x_1)$. In this connection one can speak about the *amplitude*, *gradient* and *square-gradient scattering* (SGS) mechanisms. Note that these terms are separately associated with each of two surfaces. Apart from this, there is a specific group of terms that are due to the inter-correlations between two surfaces. In what follows we refer to these terms depending on the product $\sigma^2\xi'_{\downarrow}(x)\xi'_{\uparrow}(x)$, as to the *gradient inter-product scattering* (GIS) mechanism.

It should be stressed that the GIS-terms, due to their proportionality to σ^2, were neglected in previous studies of the transport properties of surface-disordered waveguides. A similar situation occurs for the SGS-terms that have been analyzed only recently in Ref. [17, 19]. As we show below, all these terms can give the main contribution in the scattering, although formally they may be treated as negligible ones.

We start our analysis with the presentation of the kernel as a sum of its average and fluctuating parts, i.e.,

$$\Xi(k_x,k'_x)=\langle\Xi(k_x,k'_x)\rangle+\widetilde{\Xi}(k_x,k'_x). \tag{3.8}$$

It can be tested that the average, $\langle\Xi(k_x,k'_x)\rangle$, contributes only to the real part γ_n of the complex renormalization δk_n of the lengthwise wave number k_n (see Eq. (1.3)), and therefore, does not change static transport properties of the surface disordered waveguide. Thus, we will omit it because our interest is in the attenuation length L_n, not in γ_n. Therefore, it is necessary to identify the terms that have a non-zero mean-value in the general expression (3.7).

IV. FLUCTUATING SURFACE SCATTERING POTENTIAL

A. Waveguide with uncorrelated boundaries

One can see that the mean value of the kernel in Eq. (3.7) is defined by the SGS-terms that are proportional to $\sigma^2{\xi'_i}^2(x)$. In order to construct the fluctuating part of $\Xi(k_x,k'_x)$, we introduce the *square-gradient functions* $\widehat{\mathcal{V}}_i(x)$ for the lower ($i=\downarrow$) and upper ($i=\uparrow$) boundary, by subtracting the mean values, $\langle{\xi'_\downarrow}^2(x)\rangle=\langle{\xi'_\uparrow}^2(x)\rangle=\langle{\xi'}^2(x)\rangle$, from ${\xi'_i}^2(x)$,

$$\widehat{\mathcal{V}}_i(x)={\xi'_i}^2(x)-\langle{\xi'}^2(x)\rangle\,,\qquad\langle\widehat{\mathcal{V}}_i(x)\rangle=0. \tag{4.1}$$

The functions $\widehat{\mathcal{V}}_i(x)$ play an important role in our further consideration. In accordance with the Gaussian nature of the surface-profile functions $\xi_i(x)$, there are no correlations between $\widehat{\mathcal{V}}_i(x)$ and both $\xi_i(x)$ and $\xi'_i(x)$,

$$\langle\xi_i(x)\widehat{\mathcal{V}}_i(x')\rangle=0\,,\qquad\langle\xi'_i(x)\widehat{\mathcal{V}}_i(x')\rangle=0\,, \tag{4.2}$$

as well as between the product $\xi'_\downarrow(x)\xi'_\uparrow(x)$ and $\widehat{\mathcal{V}}_i(x)$, $\xi_i(x)$ or $\xi'_i(x)$.

The binary correlators for $\widehat{\mathcal{V}}_i(x)$ are given by

$$\langle\widehat{\mathcal{V}}_i(x)\widehat{\mathcal{V}}_i(x')\rangle=2\langle\xi'_i(x)\xi'_i(x')\rangle^2=2{\mathcal{W}''}^2(x-x'). \tag{4.3}$$

Correspondingly, the Fourier transform of $\widehat{\mathcal{V}}_i(x)$ reads,

$$\langle V_i(k_x)V_i(k'_x)\rangle=4\pi\delta(k_x+k'_x)\,T(k_x)\,, \tag{4.4}$$

due to the standard relation,

$$V_i(k_x)\;=\;\int_{-\infty}^{\infty}dx\exp(-ik_xx)\widehat{\mathcal{V}}_i(x). \tag{4.5}$$

Here we have introduced the *roughness-square-gradient power* (RSGP) spectrum $T(k_x)$,

$$T(k_x)=\int_{-\infty}^{\infty}dx\exp{(-ik_xx)}{\mathcal{W}''}^2(x). \tag{4.6}$$

We also introduce the Fourier transform of the product $\xi'_\downarrow(x)\xi'_\uparrow(x)$ and its binary correlator ,

$$V_\updownarrow(k_x)=\int_{-\infty}^{\infty}dx\exp(-ik_xx)\xi'_\downarrow(x)\xi'_\uparrow(x), \tag{4.7}$$

$$\langle V_\updownarrow(k_x)V_\updownarrow(k'_x)\rangle=2\pi\delta(k_x+k'_x)\,T(k_x), \tag{4.8}$$

as well as the Fourier transform of $\xi_i(x)$,

$$\widetilde{\xi}_i(k_x)\;=\;\int_{-\infty}^{\infty}dx\;\exp(-ik_xx)\,\xi_i(x). \tag{4.9}$$

It should be pointed out that by the integration by parts the power spectrum of the roughness gradients $\xi'_i(x)$ can be reduced to the RHP spectrum $W(k_x)$. However it is not possible to do for the RSGP spectrum $T(k_x)$. This very fact reflects a highly non-trivial role of the SGS- and GIS-terms, since they can compete with the other terms, although they are proportional to σ^2.

We can write now the fluctuating part of the total scattering potential in the form,

$$\begin{aligned}\widetilde{\Xi}(k_x,k'_x)&=\Xi_{1\uparrow}(k_x,k'_x)+\Xi_{1\downarrow}(k_x,k'_x)\\&+\Xi_{2\uparrow}(k_x,k'_x)+\Xi_{2\downarrow}(k_x,k'_x)+\Xi_{2\updownarrow}(k_x,k'_x).\end{aligned} \tag{4.10}$$

Here the first term is associated with the terms depending in Eq. (3.7) on $\sigma\xi_\uparrow(x)$ and $\sigma\xi'_\uparrow(x)$, therefore, on the amplitude and gradient scattering mechanisms, respectively,

$$\begin{aligned}&\Xi_{1\uparrow}(k_x,k'_x)=\\&\frac{\sigma}{d}\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\,\frac{\sin(k_zz_1)}{k_z}\\&\times\left\{2\xi_\uparrow(x_1)\frac{\partial^2}{\partial z_1^2}+i(k_x+k'_x)\xi'_\uparrow(x_1)\right.\\&\left.\times\left[\frac{1}{2}+z_1\frac{\partial}{\partial z_1}\right]\right\}\exp(ik'_xx_1)\,\frac{\sin(k'_zz_1)}{k'_z}\end{aligned} \tag{4.11a}$$

$$\begin{aligned}&\Xi_{1\uparrow}(k_n,k_{n'})=\\&-\sigma\{\delta_{nn'}+\cos[\pi(n-n')](1-\delta_{nn'})\}\tilde{\xi}_\uparrow(k_n-k_{n'}).\end{aligned} \tag{4.11b}$$

The second term in Eq. (4.10) corresponding to the lower boundary, is of the same form,

$$\begin{aligned}&\Xi_{1\downarrow}(k_x,k'_x)=\\&-\frac{\sigma}{d}\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\,\frac{\sin(k_zz_1)}{k_z}\end{aligned}$$

$$\times \left\{ 2\xi_{\downarrow}(x_1)\frac{\partial^2}{\partial z_1^2} + i(k_x + k'_x)\xi'_{\downarrow}(x_1) \right.$$
$$\left. \times \left[\frac{1}{2} + (z_1 - d)\,\frac{\partial}{\partial z_1}\right]\right\} \exp(ik'_x x_1)\,\frac{\sin(k'_z z_1)}{k'_z} \quad (4.12a)$$

$$\Xi_{1\downarrow}(k_n, k_{n'}) = \sigma\,\tilde{\xi}_{\downarrow}(k_n - k_{n'}). \quad (4.12b)$$

Next two terms in Eq. (4.10) are the SGS-terms, with ${\xi'_i}^2(x)$ being replaced by $\hat{\mathcal{V}}_i(x)$,

$$\Xi_{2\uparrow}(k_x, k'_x) =$$
$$-\frac{\sigma^2}{d^2}\int_{-\infty}^{\infty} dx_1 \int_0^d dz_1 \exp(-ik_x x_1)\,\frac{\sin(k_z z_1)}{k_z}$$
$$\times \hat{\mathcal{V}}_{\uparrow}(x_1)\left[\frac{1}{4} + 2z_1\,\frac{\partial}{\partial z_1} + z_1^2\frac{\partial^2}{\partial z_1^2}\right]$$
$$\times \exp(ik'_x x_1)\,\frac{\sin(k'_z z_1)}{k'_z} \quad (4.13a)$$

$$\Xi_{2\uparrow}(k_n, k_{n'}) =$$
$$\frac{\sigma^2 d}{2}\left\{\left[\frac{1}{3} + \frac{1}{(2\pi n)^2}\right]\delta_{nn'} + \frac{4}{\pi^2}\,\frac{n^2 + n'^2}{(n^2 - n'^2)^2}\right.$$
$$\left. \times \cos[\pi(n - n')]\,(1 - \delta_{nn'})\right\} V_{\uparrow}(k_n - k_{n'}). \quad (4.13b)$$

$$\Xi_{2\downarrow}(k_x, k'_x) =$$
$$-\frac{\sigma^2}{d^2}\int_{-\infty}^{\infty} dx_1 \int_0^d dz_1 \exp(-ik_x x_1)\,\frac{\sin(k_z z_1)}{k_z}$$
$$\times \hat{\mathcal{V}}_{\downarrow}(x_1)\left[\frac{1}{4} + 2\,(z_1 - d)\,\frac{\partial}{\partial z_1} + (z_1 - d)^2\,\frac{\partial^2}{\partial z_1^2}\right]$$
$$\times \exp(ik'_x x_1)\,\frac{\sin(k'_z z_1)}{k'_z} \quad (4.14a)$$

$$\Xi_{2\downarrow}(k_n, k_{n'}) =$$
$$\frac{\sigma^2 d}{2}\left\{\left[\frac{1}{3} + \frac{1}{(2\pi n)^2}\right]\delta_{nn'}\right.$$
$$\left. + \frac{4}{\pi^2}\,\frac{n^2 + n'^2}{(n^2 - n'^2)^2}\,(1 - \delta_{nn'})\right\} V_{\downarrow}(k_n - k_{n'}). \quad (4.14b)$$

Finally, the last term in Eq. (4.10) is due to the GIS terms in Eq. (3.7),

$$\Xi_{2\updownarrow}(k_x, k'_x) =$$
$$\frac{\sigma^2}{d^2}\int_{-\infty}^{\infty} dx_1 \int_0^d dz_1 \exp(-ik_x x_1)\,\frac{\sin(k_z z_1)}{k_z}$$
$$\times \xi'_{\downarrow}(x_1)\xi'_{\uparrow}(x_1)\left[\frac{1}{2} + 2(2z_1 - d)\,\frac{\partial}{\partial z_1} + 2z_1(z_1 - d)\frac{\partial^2}{\partial z_1^2}\right]$$
$$\times \exp(ik'_x x_1)\,\frac{\sin(k'_z z_1)}{k'_z} \quad (4.15a)$$

$$\Xi_{2\updownarrow}(k_n, k_{n'}) =$$
$$\frac{\sigma^2 d}{2}\left\{\left[\frac{1}{3} - \frac{1}{2(\pi n)^2}\right]\delta_{nn'} - \frac{8}{\pi^2}\,\frac{n^2 + n'^2}{(n^2 - n'^2)^2}\right.$$
$$\left. \times \cos^2[\pi(n - n')/2]\,(1 - \delta_{nn'})\right\} V_{\updownarrow}(k_n - k_{n'}). \quad (4.15b)$$

The kernel $\widetilde{\Xi}(k_x, k'_x)$ written as the sum (4.10) of specially designed terms, has a very convenient form. First, all the terms are chosen to have zero average. Second, since the functions $\xi_i(x)$, $\xi_{\downarrow}(x)\xi_{\uparrow}(x)$ and $\widehat{\mathcal{V}}_i(x)$ do not correlate with each other, their Fourier transforms are also non-correlated:

$$\langle\tilde{\xi}_i(k_x)V_i(k'_x)\rangle = 0. \quad (4.16a)$$
$$\langle\tilde{\xi}_i(k_x)V_{\updownarrow}(k'_x)\rangle = 0. \quad (4.16b)$$
$$\langle V_i(k'_x)V_{\updownarrow}(k_x)\rangle = 0. \quad (4.16c)$$

Therefore, all the terms in Eq. (4.10) determining the potential, are also non-correlated. Their autocorrelators are,

$$\langle\Xi_{1i}(k_n, k_{n'})\Xi_{1i}(k_{n'}, k_{n''})\rangle = 2\pi\delta(k_n - k_{n''})Q_1(k_n, k_{n'}), \quad (4.17a)$$

$$Q_1(k_n, k_{n'}) = \sigma^2 W(k_n - k_{n'}), \quad (4.17b)$$

$$\langle\Xi_{2i}(k_n, k_{n'})\Xi_{2i}(k_{n'}, k_{n''})\rangle = 2\pi\delta(k_n - k_{n''})Q_2(k_n, k_{n'}), \quad (4.18a)$$

$$Q_2(k_n, k_{n'}) = \frac{\sigma^4 d^2}{2}T(k_n - k_{n'})\left\{\left[\frac{1}{3} + \frac{1}{(2\pi n)^2}\right]^2\delta_{nn'}\right.$$
$$\left. + \frac{16}{\pi^4}\,\frac{(n^2 + n'^2)^2}{(n^2 - n'^2)^4}\,(1 - \delta_{nn'})\right\}, \quad (4.18b)$$

$$\langle\Xi_{2\updownarrow}(k_n, k_{n'})\Xi_{2\updownarrow}(k_{n'}, k_{n''})\rangle = 2\pi\delta(k_n - k_{n''})Q_{2\updownarrow}(k_n, k_{n'}) \quad (4.19a)$$

$$Q_{2\updownarrow}(k_n, k_{n'}) = \frac{\sigma^4 d^2}{4}T(k_n - k_{n'})\left\{\left[\frac{1}{3} - \frac{1}{2(\pi n)^2}\right]^2\delta_{nn'}\right.$$
$$\left. + \frac{64}{\pi^4}\,\frac{(n^2 + n'^2)^2}{(n^2 - n'^2)^4}\,\cos^4[\pi(n - n')/2]\,(1 - \delta_{nn'})\right\}. \quad (4.19b)$$

Note that in spite of the absence of correlations between the upper and lower surface profiles, there are the correlations due to their derivatives since $\langle\Xi_{2\updownarrow}(k_n, k_{n'})\Xi_{2\updownarrow}(k_{n'}, k_{n''})\rangle \neq 0$ (see Eq. (4.19a)).

As a result, the correlator of the fluctuating scattering potential (4.10) in the k_x-representation reads,

$$\langle\widetilde{\Xi}(k_n,k_{n'})\widetilde{\Xi}(k_{n'},k_{n''})\rangle = 2\pi\delta(k_n-k_{n''})Q(k_n,k_{n'}), \tag{4.20a}$$

$$Q(k_n,k_{n'}) = 2Q_1(k_n,k_{n'})+2Q_2(k_n,k_{n'})+Q_{2\updownarrow}(k_n,k_{n'})\,. \tag{4.20b}$$

The factor 2 in the first and second terms arises due to independent contributions, $Q_1(k_n,k_{n'})$ and $Q_2(k_n,k_{n'})$, from the lower and upper boundaries.

B. Waveguide with antisymmetric rough boundaries

In this configuration both boundaries of the waveguide are identical, see Eq. (1.1), therefore, antisymmetric with respect to the central line $z=d/2$. For this specific case the width of the waveguide (the distance between upper and lower boundaries) remains constant along the $x-$direction. Instead of Eq. (3.7) we have to start now with the exact expression ((3.6)) by substituting the relation ((1.1)). As a result, one gets the relatively simple expression for the scattering potential,

$$\begin{aligned}\Xi(k_x,k'_x) &= \int_{-\infty}^{\infty}dx_1\int_0^d dz_1\,\exp\left(-ik_xx_1\right)\frac{\sin(k_zz_1)}{k_z}\\ &\times\left(\sigma\left[i(k_x+k'_x)\right]\xi'(x_1)\,\frac{\partial}{\partial z_1}\right.\\ &\left.-\,\sigma^2\,{\xi'}^2(x_1)\frac{\partial^2}{\partial {z_1}^2}\right)\exp\left(ik'_xx_1\right)\frac{\sin(k'_zz_1)}{k'_z}\end{aligned} \tag{4.21}$$

One can realize that all terms proportional to σ^2 can be written as one SGS-term with a non-zero mean value. By making use of the functions defined by Eqs. (4.1), (4.5) and (4.9) with $i=\uparrow$, in what follows we omit the subscript i.

Now the fluctuating part of the scattering potential is,

$$\widetilde{\Xi}(k_x,k'_x)=\Xi_1(k_x,k'_x)+\Xi_2(k_x,k'_x). \tag{4.22}$$

The first summand related to the gradient scattering terms in Eq. (4.21) has the form,

$$\begin{aligned}\Xi_1(k_x,k'_x) &= \sigma\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\frac{\sin(k_zz_1)}{k_z}\\ &\times[i(k_x+k'_x)]\,\xi'(x_1)\,\frac{\partial}{\partial z_1}\exp(ik'_xx_1)\frac{\sin(k'_zz_1)}{k'_z}\end{aligned} \tag{4.23a}$$

$$\Xi_1(k_n,k_{n'})=2\sigma\sin^2[\pi(n-n')/2]\,\tilde{\xi}(k_n-k_{n'}). \tag{4.23b}$$

The second summand is due to the SGS-term, however, with the function $\hat{\mathcal{V}}(x_1)$ instead of ${\xi'}^2(x_1)$. With the use of the Fourier transform (4.5) for the operator $\hat{\mathcal{V}}(x)$, one gets,

$$\begin{aligned}\Xi_2(k_x,k'_x) &= -\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\frac{\sin(k_zz_1)}{k_z}\\ &\times\sigma^2\hat{\mathcal{V}}(x_1)\,\frac{\partial^2}{\partial z_1^2}\exp(ik'_xx_1)\frac{\sin(k'_zz_1)}{k'_z}\end{aligned} \tag{4.24a}$$

$$\Xi_2(k_n,k_{n'})=\frac{\sigma^2d}{2}\,\delta_{nn'}\,V(k_n-k_{n'}). \tag{4.24b}$$

Note that the expressions (4.23) and (4.24) for the scattering potential are still *exact*. In contrast with other configurations discussed below, the expansion in σ/d-smallness is not required because of the constant width of the waveguide.

The correlation properties of the scattering potential are described by the following formulas. Due to the condition (4.16a), the terms $\Xi_1(k_x,k'_x)$ and $\Xi_2(k_x,k'_x)$ are non-correlated,

$$\langle\Xi_1(k_x,k'_x)\Xi_2(k'_x,k''_x)\rangle=0. \tag{4.25}$$

On the other hand, the autocorrelators are non-zero,

$$\langle\Xi_1(k_n,k_{n'})\Xi_1(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q_1(k_n,k_{n'}), \tag{4.26a}$$

$$Q_1(k_n,k_{n'})=4\sigma^2W(k_n-k_{n'})\sin^4[\pi(n-n')/2]. \tag{4.26b}$$

$$\langle\Xi_2(k_n,k_{n'})\Xi_2(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q_2(k_n,k_{n'}), \tag{4.27a}$$

$$Q_2(k_n,k_{n'})=(\sigma^4d^2/2)\,T(k_n-k_{n'})\,\delta_{nn'}\,. \tag{4.27b}$$

Finally, the correlator of the fluctuating part of the total scattering potential $\widetilde{\Xi}(k_x,k'_x)$ has the following form,

$$\langle\widetilde{\Xi}(k_n,k_{n'})\widetilde{\Xi}(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q(k_n,k_{n'}), \tag{4.28a}$$

$$Q(k_n,k_{n'})=Q_1(k_n,k_{n'})+Q_2(k_n,k_{n'}). \tag{4.28b}$$

C. Waveguide with symmetric rough boundaries

Let us now consider the waveguide with symmetric boundaries, see Eq. (1.2), therefore, with the width defined by

$$w(x)=d+2\sigma\xi(x),\qquad\langle w(x)\rangle=d. \tag{4.29}$$

By substituting the symmetric profile into Eq. (3.7), we obtain for the surface scattering potential,

$$\Xi(k_x,k'_x)\approx\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\,\exp\left(-ik_xx_1\right)\frac{\sin(k_zz_1)}{k_z}$$

$$\times\left\{\frac{4\sigma\xi(x_1)}{d}\frac{\partial^2}{\partial z_1^2}\right.$$

$$+i(k_x+k'_x)\frac{\sigma\xi'(x_1)}{d}\left[1+(2z_1-d)\,\frac{\partial}{\partial z_1}\right]$$

$$\left.-\,\frac{\sigma^2\xi'^2(x_1)}{d^2}\left[1+4\,(2z_1-d)\,\frac{\partial}{\partial z_1}+(2z_1-d)^2\,\frac{\partial^2}{\partial z_1^2}\right]\right\}$$

$$\times\exp\left(ik'_xx_1\right)\frac{\sin(k'_zz_1)}{k'_z}\,. \tag{4.30}$$

As for the previous configuration, see Eq. (4.21), the SGS-terms corresponding to the lower boundary and the GIS-terms are written in terms of $\xi(x)=\xi_\uparrow(x)$.

The potential in Eq. (4.30) can be written as the sum of two terms related to substantially different scattering mechanisms,

$$\widetilde{\Xi}(k_x,k'_x)=\Xi_1(k_x,k'_x)+\Xi_2(k_x,k'_x). \tag{4.31}$$

Here the first summand is related to the terms proportional to σ in Eq. (4.30) (the amplitude and gradient scattering terms),

$$\Xi_1(k_x,k'_x)=\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\,\frac{\sin(k_zz_1)}{k_z}$$

$$\times\left\{\frac{4\sigma\xi(x_1)}{d}\frac{\partial^2}{\partial z_1^2}+i(k_x+k'_x)\frac{\sigma\xi'(x_1)}{d}\right.$$

$$\left.\times\left[1+(2z_1-d)\,\frac{\partial}{\partial z_1}\right]\right\}\exp(ik'_xx_1)\,\frac{\sin(k'_zz_1)}{k'_z} \tag{4.32a}$$

$$\Xi_1(k_n,k_{n'})=-2\sigma\{\delta_{nn'}+\cos^2[\pi(n-n')/2]$$

$$\times(1-\delta_{nn'})\}\,\tilde{\xi}(k_n-k_{n'}). \tag{4.32b}$$

The second summand is associated with the fluctuating part of the SGS-terms in Eq. (4.30), and consequently, with the function ${\xi'}^2(x)$ being replaced by $\hat{\mathcal{V}}(x)$,

$$\Xi_2(k_x,k'_x)=-\int_{-\infty}^{\infty}dx_1\int_0^d dz_1\exp(-ik_xx_1)\,\frac{\sin(k_zz_1)}{k_z}$$

$$\times\frac{\sigma^2\hat{\mathcal{V}}(x_1)}{d^2}\left[1+4\,(2z_1-d)\,\frac{\partial}{\partial z_1}+(2z_1-d)^2\,\frac{\partial^2}{\partial z_1^2}\right]$$

$$\times\exp(ik'_xx_1)\,\frac{\sin(k'_zz_1)}{k'_z} \tag{4.33a}$$

$$\Xi_2(k_n,k_{n'})=\frac{\sigma^2d}{2}\left\{\left[\frac{1}{3}+\frac{1}{(\pi n)^2}\right]\delta_{nn'}+\frac{16}{\pi^2}\right.$$

$$\left.\times\frac{n^2+n'^2}{(n^2-n'^2)^2}\cos^2[\pi(n-n')/2](1-\delta_{nn'})\right\}V(k_n-k_{n'}). \tag{4.33b}$$

The correlation properties of the introduced surface scattering potentials $\Xi_1(k_n,k_{n'})$ and $\Xi_2(k_n,k_{n'})$ are described by the correlator,

$$\langle\Xi_1(k_n,k_{n'})\Xi_1(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q_1(k_n,k_{n'}), \tag{4.34a}$$

$$Q_1(k_n,k_{n'})=4\sigma^2W(k_n-k_{n'})$$

$$\times\{\delta_{nn'}+\cos^4[\pi(n-n')/2](1-\delta_{nn'})\}. \tag{4.34b}$$

$$\langle\Xi_2(k_n,k_{n'})\Xi_2(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q_2(k_n,k_{n'}), \tag{4.35a}$$

$$Q_2(k_n,k_{n'})=\frac{\sigma^4d^2}{2}T(k_n-k'_x)\left\{\left[\frac{1}{3}+\frac{1}{(\pi n)^2}\right]^2\delta_{nn'}\right.$$

$$\left.+\frac{256}{\pi^4}\,\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4}\cos^4[\pi(n-n')/2](1-\delta_{nn'})\right\}. \tag{4.35b}$$

Thus, we arrive at the expression,

$$\langle\widetilde{\Xi}(k_n,k_{n'})\widetilde{\Xi}(k_{n'},k_{n''})\rangle=2\pi\delta(k_n-k_{n''})Q(k_n,k_{n'}), \tag{4.36a}$$

$$Q(k_n,k_{n'})=Q_1(k_n,k_{n'})+Q_2(k_n,k_{n'}). \tag{4.36b}$$

V. AVERAGE GREEN'S FUNCTION AND ATTENUATION LENGTH

Now we replace the problem for the random Green's function $g(k_x,k'_x)$ with the problem for the Green's function $\langle g(k_x,k'_x)\rangle$ averaged over the surface disorder. To perform the averaging of Eq. (3.1) with $\Xi(k_x,k'_x)$, given for the uncorrelated, antisymmetric and symmetric boundary configurations by Eq. (4.10), (4.22) and (4.31), respectively, and obtain $\langle g(k_x,k'_x)\rangle$, we can apply one of the standard and well known perturbative methods. For example, it can be the diagrammatic approach developed for surface disordered systems [1], as well as the technique developed in Ref. 3. Both of the methods take adequately into account the multiple scattering from the corrugated boundaries, and allow one to develop the consistent perturbative approach with respect to the scattering potential. After quite cumbersome calculations (see Ref. [17] for details), we obtain the average Green's function in the normal-mode representation,

$$\langle\mathcal{G}(x,x';z,z')\rangle=\sum_{n=1}^{N_d}\sin\left(\frac{\pi nz}{d}\right)\sin\left(\frac{\pi nz'}{d}\right)$$

$$\times\frac{\exp(ik_n|x-x'|)}{ik_nd}\,\exp\left(-\frac{|x-x'|}{2L_n}\right). \tag{5.1}$$

The quantity L_n is the wave attenuation length or electron total mean-free-path of the nth conducting mode. This quantity is determined by the imaginary part of the self-energy and describes the scattering from the nth mode into other propagating modes,

$$\frac{1}{L_n} = \frac{(\pi n/d)^2}{k_n d} \sum_{n'=1}^{N_d} \frac{(\pi n'/d)^2}{k_{n'} d} \times \left[Q(k_n, -k_{n'}) + Q(k_n, +k_{n'}) \right]. \qquad (5.2)$$

Here the explicit expressions for $Q(k_n, -k_{n'})$ and $Q(k_n, +k_{n'})$ are given by Eqs. (4.20b), (4.28b) and (4.36b) for the waveguide with uncorrelated, antisymmetric and symmetric boundaries.

In deriving Eqs. (5.1) and (5.2) we have made the following simplifications. First, the self-energy in the Dyson-type equation for the average Green's function was obtained within the second-order approximation in the perturbation potential. In terms of the diagrammatic technique this is similar to the "simple vortex" or, the same, Bourret approximation [20] which contains the binary correlator $Q(k_x, k_x')$ of the surface scattering potential and the unperturbed pole factor $g_0(q_x)$. Second, in order to extract the inverse attenuation length from the self-energy, we substituted the lengthwise wave number k_x by its unperturbed value k_n. The justification of this substitution, as well as the conditions of applicability resulting from the above simplifications, were discussed in Ref. [17]. Specifically, it was shown that the domain of applicability is restricted by two independent criteria of the *weak surface scattering*,

$$\Lambda_n = k_n d/(\pi n/d) \ll 2L_n, \qquad R_c \ll 2L_n. \qquad (5.3)$$

Here Λ_n is the distance between two successive reflections of the nth mode from one boundary to the opposite one. The conditions in (5.3) imply that the wave is weakly attenuated on both the correlation length R_c and the cycle length Λ_n. They implicitly include the requirement that $\sigma \ll d$ for the surface corrugations be small in height, used in the Sec. IV when deriving the explicit form (3.7) for the surface scattering potential $\Xi(k_x, k_x')$.

VI. ATTENUATION LENGTH ANALYSIS

In this section we discuss the attenuation length L_n for three configurations of the waveguide. In the first part of the subsections we give explicit expressions for the $1/L_n$ for any form of the binary correlator $\mathcal{W}(x)$. In the second part, we assume that the random surface profile $\xi(x)$ has the Gaussian binary correlator,

$$\mathcal{W}(x) = \exp(-x^2/2R_c^2). \qquad (6.1)$$

Then the RHP spectrum (2.7) is given by

$$W(k_x) = \sqrt{2\pi}\, R_c \, \exp(-k_x^2 R_c^2/2). \qquad (6.2)$$

The RSGP spectrum $T(k_x)$ defined by Eq. (4.6)), can be written as

$$T(k_x) = \frac{\sqrt{\pi}}{16 R_c^3} \left[(k_x R_c)^4 - 4(k_x R_c)^2 + 12 \right] \times \exp\left[-(k_x R_c)^2/4 \right]. \qquad (6.3)$$

In what follows, we express the inverse attenuation length in the form of the dimensionless quantity $\Lambda_n/2L_n$. Since L_n depends on as many as four dimensionless parameters $(k\sigma)^2$, kR_c, kd/π, and n, a complete analysis appears to be quite complicated. For this reason, below we restrict ourselves to an analysis of the interplay between different scattering mechanisms, as a function of the dimensionless correlation length kR_c for different values of $(k\sigma)^2$. We concentrate our attention on a multimode waveguide with a large number of propagating modes, $N_d \approx kd/\pi \gg 1$. Our main interest is in two situations, namely, in a small-scale roughness of the "white noise" kind for which $kR_c \ll 1$, and in large-scale random corrugations with $kR_c \gg 1$.

A. Waveguide with uncorrelated boundaries

In view of Eq. (4.20b), the general expression (5.2) for the inverse attenuation length shows that for a waveguide with uncorrelated boundaries this quantity consists of three terms,

$$\frac{1}{L_n} = \frac{1}{L_n^{(1)}} + \frac{1}{L_n^{(2)}} + \frac{1}{L_n^{(2)\updownarrow}}. \qquad (6.4)$$

These terms are originated from different mechanisms of the surface scattering. The first attenuation length $L_n^{(1)}$ is related to the RHP spectrum $W(k_x)$ through the expression for $2Q_1(k_x, k_x')$. In accordance with Eq. (4.17b) it is given by

$$\frac{1}{L_n^{(1)}} = 2\sigma^2 \frac{(\pi n/d)^2}{k_n d} \sum_{n'=1}^{N_d} \frac{(\pi n'/d)^2}{k_{n'} d} \times \left[W(k_n + k_{n'}) + W(k_n - k_{n'}) \right]. \qquad (6.5)$$

Here the term $1/L_{n,n}^{(1)}$ corresponding to $n' = n$, is related to the amplitude scattering while the terms $1/L_{n,n'\neq n}^{(1)}$ result from the gradient scattering. These two mechanisms of surface scattering are due to the corresponding terms in the expression for $\Xi_{1i}(k_x, k_x')$, see Eqs. (4.11a) and (4.12a). The first one is related to the term depending on the amplitude of the roughness profile $\sigma\xi_i(x)$, and the second one is due to the terms depending on the roughness gradient $\sigma\xi_i'(x)$. The expression (6.5) exactly coincides with that previously obtained by various methods (see, e.g., Ref. 1).

The second attenuation length $L_n^{(2)}$ related to the RSGP spectrum through $2Q_2(k_x, k_x')$, is associated solely with the SGS mechanism due to the operator $\widehat{\mathcal{V}}_i(x)$, see Eqs. (4.13a) and (4.14a). In accordance with Eq. (4.18b), it is described by

$$\frac{1}{L_n^{(2)}} = \sum_{n'=1}^{N_d} \frac{1}{L_{n,n'}^{(2)}}. \qquad (6.6)$$

Here the terms in the sum are

$$\frac{1}{L^{(2)}_{n,n}} = \sigma^4 \frac{(\pi n/d)^4}{k_n^2}\left[\frac{1}{3}+\frac{1}{(2\pi n)^2}\right]^2 [T(2k_n)+T(0)] \tag{6.7}$$

and

$$\begin{aligned}\frac{1}{L^{(2)}_{n,n'\neq n}} =&\sigma^4 \frac{16}{\pi^4}\frac{(\pi n/d)^2}{k_n}\frac{(\pi n'/d)^2}{k_{n'}}\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4} \\ &\times \left[T(k_n+k_{n'})+T(k_n-k_{n'})\right].\end{aligned} \tag{6.8}$$

The third term $L_n^{(2)\updownarrow}$ is also related to the RSGP spectrum through $Q_{2\updownarrow}(k_x,k'_x)$ due to the product $\xi'_{\downarrow}(x)\xi'_{\uparrow}(x)$, see Eq. (4.15a). In accordance with Eqs. (4.19b), it is described by

$$\frac{1}{L_n^{(2)\updownarrow}} = \sum_{n'=1}^{N_d}\frac{1}{L^{(2)\updownarrow}_{n,n'}}, \tag{6.9}$$

where

$$\begin{aligned}\frac{1}{L^{(2)\updownarrow}_{n,n}} =&\frac{\sigma^4}{4}\frac{(\pi n/d)^4}{k_n^2}\left[\frac{1}{3}-\frac{1}{2(\pi n)^2}\right]^2 \\ &\times \left[T(2k_n)+T(0)\right].\end{aligned} \tag{6.10}$$

and

$$\begin{aligned}\frac{1}{L^{(2)\updownarrow}_{n,n'\neq n}} =& \sigma^4 \frac{16}{\pi^4}\frac{(\pi n/d)^2}{k_n}\frac{(\pi n'/d)^2}{k_{n'}}\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4} \\ &\times \left[T(k_n+k_{n'})+T(k_n-k_{n'})\right]\cos^4[\pi(n-n')/2].\end{aligned} \tag{6.11}$$

Analogously, the terms in Eq. (6.4) can be classified as partial terms describing the intramode ($1/L^{(1)}_{n,n}$, $1/L^{(2)}_{n,n}$ and $1/L^{(2)\updownarrow}_{n,n}$), and intermode ($1/L^{(1)}_{n,n'\neq n}$, $1/L^{(2)}_{n,n'\neq n}$ and $1/L^{(2)\updownarrow}_{n,n'\neq n}$) scattering. According to the discussion of Eq. (6.5), the amplitude and gradient terms result in the intramode and intermode scattering, respectively. As for the square-gradient terms, they lead to both types of scattering.

1. Small-scale roughness

To continue our analysis, let us start with the widely used case of *a small-scale boundary perturbation*, $kR_c \ll 1$. In this case the surface roughness can be regarded as a delta-correlated random process with the correlator $\mathcal{W}(x-x') \approx W(0)\delta(x-x')$ and constant power spectrum $W(k_x) \approx W(0) \sim R_c$. Taking into account the evident relationship $k\Lambda_n \gtrsim 1$, one can get the following inequalities specifying this case,

$$kR_c \ll 1 \lesssim k\Lambda_n. \tag{6.12}$$

It is necessary to underline that in the regime of small-scale roughness (6.12) the second of the weak-scattering conditions in Eq.(5.3) is not so restrictive as the first one, and directly stems from it, $R_c \ll \Lambda_n \ll 2L_n$.

In Eqs. (6.5), (6.7), (6.8), (6.10) and (6.11) for the attenuation lengths the argument of the correlators $W(k_x)$ and $T(k_x)$ turns out to be much less than the scale of their decrease R_c^{-1} under the conditions (6.12). Thus, for any term in the summation over n' the argument can be taken as zero.

Therefore, the first attenuation length is determined as follows,

$$\frac{\Lambda_n}{2L_n^{(1)}} \approx 2(k\sigma)^2\frac{n}{kd/\pi}\frac{W(0)}{k}\sum_{n'=1}^{N_d}\frac{(\pi n'/d)^2}{k_{n'}d} \tag{6.13a}$$

$$\approx (k\sigma)^2\frac{n}{kd/\pi}\frac{kW(0)}{2} \tag{6.13b}$$

The approximation is made here for a large number of the conducting modes, with the change of the summation over n' by integration. For a further estimate, one can take into account the formula

$$W(0) = \int_{-\infty}^{\infty} dx\mathcal{W}(x) = 2R_c\int_0^{\infty} d\rho\mathcal{W}(R_c\rho), \tag{6.14}$$

that directly follows from the definition (2.7) for the Fourier transform $W(k_x)$ of the binary correlator $\mathcal{W}(x)$. The function $\mathcal{W}(R_c\rho)$ is the dimensionless correlator of the dimensionless variable ρ, with the scale of decrease of the order of 1. As a result, the function $\mathcal{W}(R_c\rho)$ does not depend on R_c. Therefore, the integral over ρ entering Eq. (6.14) is a positive constant of the order of unity. For example, in the case of the Gaussian correlations (see Eq. (6.2)), we have, $W(0) = \sqrt{2\pi}R_c$, therefore, the integral is $\sqrt{\pi/2}$.

For the second length in Eq. (6.4) one gets,

$$\begin{aligned}\frac{\Lambda_n}{2L_n^{(2)}} \approx& \pi^2\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3} \\ &\times\left\{\left[\frac{1}{3}+\frac{1}{(2\pi n)^2}\right]^2+\frac{16k_nd}{\pi^4(\pi n/d)^2}\right. \\ &\times\left.\left(\sum_{n'=1}^{n-1}+\sum_{n'=n+1}^{N_d}\right)\frac{(\pi n'/d)^2}{k_{n'}d}\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4}\right\}\end{aligned} \tag{6.15a}$$

$$\approx \frac{\pi^2}{4}\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}. \tag{6.15b}$$

In Eq. (6.15a) every term in the sum rapidly decreases with an increase of the absolute value of $\Delta n = n-n'$. This can be seen by making use of the following estimate,

$$\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4} \approx \frac{1}{4(\Delta n)^4} \qquad \text{for} \quad n \gg |\Delta n|. \tag{6.16}$$

Therefore, the sum in Eq. (6.15a) can be correctly estimated by three terms with $n' = n, n \pm 1$. For simplicity, in Eq. (6.15b) we assume $N_d \gg n \gg 1$, and replace the curly braces by factor 1/4.

The explicit form for $T(0)$ directly follows from the definition (4.6) for the correlator $T(k_x)$,

$$\begin{aligned} T(0) &= \int_{-\infty}^{\infty} dx \mathcal{W}''^2(x) \\ &= R_c^{-3} \int_{-\infty}^{\infty} d\rho \left[\frac{d^2 \mathcal{W}(R_c\rho)}{d\rho^2} \right]^2 . \end{aligned} \tag{6.17}$$

If the roughness correlations are of the Gaussian form, then according to Eq. (6.3), we have $T(0) = 3\sqrt{\pi}/4R_c^3$ and the integral over ρ entering Eq. (6.17) is equal to $3\sqrt{\pi}/4$.

The second term in Eq. (6.4) is bigger than the third one. Therefore, a good aproximation for Eq. (6.4) is

$$\frac{1}{L_n} \approx \frac{1}{L_n^{(1)}} + \frac{1}{L_n^{(2)}}. \tag{6.18}$$

One can make sure of this if we consider in Eq. (6.11) the product of the fast decreasing factor (6.16) and $\cos^4[\pi(n-n')/2]$ (the latter cancels the terms with $\Delta n = 1, 3, ..$). Thus, we arrive at the following estimate,

$$\frac{\Lambda_n}{2L_n^{(2)\updownarrow}} \approx \frac{\Lambda_n}{2L_{n,n}^{(2)\updownarrow}} \approx \frac{\pi^2}{36} \frac{(k\sigma)^4}{k_n d} \frac{n^3}{(kd/\pi)} \frac{T(0)}{k^3}. \tag{6.19}$$

By comparing Eq. (6.19) with Eq. (6.15b) one can confirm Eq. (6.18).

The expression (6.18) works well in the whole domain of applicability of our theory, not only in the region of small-scale roughness. To support this, in Fig. 1 we compare the plot of $\Lambda_n/2L_n^{(2)}$ with the plot of $\Lambda_n/2L_n^{(2)} + \Lambda_n/2L_n^{(2)\updownarrow}$; they are practically the same. Therefore, in what follows for this case of uncorrelated boundaries we will neglect the third term in Eq. (6.4).

Due to Eq. (6.18), the relationship between the first and second attenuation lengths is expressed by

$$\frac{L_n^{(1)}}{L_n^{(2)}} \sim (k\sigma)^2 \frac{n^2}{k_n d} (kR_c)^{-4}. \tag{6.20}$$

According to a substantially different behavior of the quantities $\Lambda_n/2L_n^{(1)}$ and $\Lambda_n/2L_n^{(2)}$ with respect to kR_c, it becomes clear that they have to intersect at the crossing point $(kR_c)_{cr}$. If the crossing point falls onto the region of a small-scale roughness ($kR_c \ll 1$), its dependence on the model parameters is obtained by equating the expression (6.20) to unity,

$$(kR_c)^2_{cr} \sim (k\sigma)n/\sqrt{k_n d}. \tag{6.21}$$

To the left from this point $(kR_c)_{cr}$ the SGS length prevails, $L_n^{(2)} \ll L_n^{(1)}$. To its right the main contribution is due to the first attenuation length, $L_n^{(1)} \ll L_n^{(2)}$. The expression (6.21) shows that the crossing point is smaller for smaller values of $k\sigma$, as well as for smaller mode indices n, or for larger values of the mode parameter kd/π.



FIG. 1: Plot of $\Lambda_n/2L_n^{(2)}$ (solid line) and $\Lambda_n/2L_n^{(2)} + \Lambda_n/2L_n^{(2)\updownarrow}$ (dashed line) versus kR_c, for $kd/\pi = 100.5$, $n = 50$ and $(k\sigma)^2 = 10^{-2}$.

2. *Large-scale roughness: weak correlations*

The intermediate situation arises when the correlation length R_c becomes much larger than the wave length $2\pi/k$, but still remains much less than the cycle length Λ_n,

$$1 \ll kR_c \ll k\Lambda_n. \tag{6.22}$$

As before, the first of the weak-scattering conditions (5.3) is the most restrictive, $R_c \ll \Lambda_n \ll 2L_n$.

Since the distance Λ_n is larger than the correlation length R_c, the successive reflections of the waves from the rough surface are weakly correlated. Note that the distance between neighboring wave numbers k_n and $k_{n\pm1}$ is much smaller than the variation scale R_c^{-1} of the correlators $W(k_x)$ and $T(k_x)$,

$$|k_n - k_{n\pm1}| \approx |\partial k_n/\partial n| = \pi\Lambda_n^{-1} \ll R_c^{-1}. \tag{6.23}$$

This implies that the correlators $W(k_x)$ and $T(k_x)$ are smooth functions of the summation index n'. Therefore, the sum in the expression (6.5) for the first attenuation length $L_n^{(1)}$ can be substituted by the integral,

$$\frac{\Lambda_n}{2L_n^{(1)}} \approx (k\sigma)^2 \frac{n}{kd/\pi} \int_0^{N_d} dn' \frac{(\pi n'/d)^2}{k_{n'} d}$$

$$\times \frac{1}{k} \left[W(k_n + k_{n'}) + W(k_n - k_{n'}) \right] \qquad (6.24a)$$

$$= \frac{(k\sigma)^2}{\pi} \frac{n}{kd/\pi} \int_{-k}^{k} dk_x \sqrt{k^2 - k_x^2} \, \frac{W(k_n - k_x)}{k}. \qquad (6.24b)$$

Eq. (6.24) shows that the first attenuation length is contributed by the scattering of a given n-th mode into other propagating modes. In order to obtain this asymptotic result, we have used only the condition of week correlations, $R_c \ll \Lambda_n$. Therefore, Eq. (6.24) provides a reduction to Eq. (6.13) for small-scale corrugations, $kR_c \ll 1$. In the case of a large-scale roughness, $kR_c \gg 1$, the formula (6.24) obtained for $L_n^{(1)}$ allows one for further simplifications as was done in Ref. 1.

In contrast with $L_n^{(1)}$, due to a rapidly decaying factor (6.16) the SGS length $L_n^{(2)}$ can be still described by keeping tree terms only, $n' = n, n \pm 1$, in the sum in Eq. (6.6). Taking into account the estimate (6.23), for the case $N_d \gg n \gg 1$ one can obtain,

$$\frac{\Lambda_n}{2L_n^{(2)}} \approx \frac{\pi^2}{2} \frac{(k\sigma)^4}{k_n d} \frac{n^3}{(kd/\pi)} \frac{T(0) + T(2k_n)}{k^3}$$

$$\times \left\{ \left[\frac{1}{3} + \frac{1}{(2\pi n)^2} \right]^2 + \frac{8}{\pi^4} \right\} \qquad (6.25a)$$

$$\approx \frac{\pi^2}{8} \frac{(k\sigma)^4}{k_n d} \frac{n^3}{(kd/\pi)} \frac{T(0) + T(2k_n)}{k^3}. \qquad (6.25b)$$

To get the final expression (6.25b), we replaced the curly braces in Eq. (6.25a) by the factor 1/4. Naturally, at small-scale corrugations, $kR_c \ll 1$, the obtained result (6.25) reduces to Eq. (6.15). For a large-scale roughness, $kR_c \gg 1$, one should use Eq. (6.25) because of an arbitrary value of the parameter $k_n R_c$. We do not consider here this case in detail due to its intermediate character.

3. *Large-scale roughness: strong correlations*

Another limit case refers to the correlation length R_c to be very large both in comparison with the wave length $2\pi/k$ and with the cycle length Λ_n,

$$1 \lesssim k\Lambda_n \ll kR_c. \qquad (6.26)$$

In this case the number of wave reflections on the scale of the correlation length R_c is large. Therefore, the successive reflections are strongly correlated to each other.

Under the relations (6.26) the second of the weak-scattering conditions (5.3) is the most restrictive, therefore, the condition of the applicability reads as

$$\Lambda_n \ll R_c \ll 2L_n. \qquad (6.27)$$

The latter requirement determines the upper limit for the value of the correlation length R_c.

Due to Eq. (6.26), the distance between neighboring wave numbers k_n and $k_{n\pm1}$ turns out to be much larger than the variation scale R_c^{-1} of the correlators $W(k_x)$ and $T(k_x)$,

$$|k_n - k_{n\pm1}| \approx |\partial k_n/\partial n| = \pi\Lambda_n^{-1} \gg R_c^{-1}. \qquad (6.28)$$

This indicates that the probability of the intermode ($n' \neq n$) transitions is exponentially small and the attenuation lengths are mainly formed by the incoherent intramode ($n' = n$) scattering. Formally, for strong correlations (6.26) the correlators $W(k_x)$ and $T(k_x)$ are sharpest functions of the summation index n'. In the sums of Eqs. (6.5) and (6.6) for the attenuation lengths the main contribution is due to the diagonal terms with $n' = n$, for which $W(2k_n)$ and $T(2k_n)$ can be neglected in comparison with $W(0)$ and $T(0)$.

Thus, the first attenuation length reads

$$\frac{\Lambda_n}{2L_n^{(1)}} \approx \frac{(k\sigma)^2}{k_n d} \frac{n^3}{(kd/\pi)^3} \, kW(0). \qquad (6.29)$$

Correspondingly, for the SGS length one gets,

$$\frac{\Lambda_n}{2L_n^{(2)}} \approx \frac{\pi^2}{2} \frac{(k\sigma)^4}{k_n d} \frac{n^3}{(kd/\pi)} \frac{T(0)}{k^3} \left[\frac{1}{3} + \frac{1}{(2\pi n)^2} \right]^2. \qquad (6.30)$$

The ratio of the first attenuation length to the second one can be presented as

$$\frac{L_n^{(1)}}{L_n^{(2)}} \sim \frac{\Lambda_n}{2L_n^{(1)}} \left(\frac{\Lambda_n}{R_c} \right)^5 (k_n \Lambda_n)^{-2} \ll 1. \qquad (6.31)$$

According to the first inequality in Eq. (5.3), applied to the condition (6.26) of the strong-correlations and to the evident relationship $k_n\Lambda_n \gtrsim 1$, one can see that the amplitude scattering length always prevails over the SGS length within the interval of strong correlations. For this reason in the condition of applicability (6.27) one should substitute L_n by $L_n^{(1)}$. This allows one to arrive at the inequalities in the explicit form,

$$k\Lambda_n \ll kR_c \ll (k\Lambda_n)(k\sigma)^{-1}(kd/\pi n). \qquad (6.32)$$

Before we pass to the next configuration of the waveguide, it is interesting to compare the case of two uncorrelated boundaries with the waveguide with one rough boundary, studied in Ref. [17]. One can found that under the approximation (6.18), the expression for $1/L_n$ turns out to be as twice as bigger than that one for the one-rough-boundary configuration. This fact is quite natural since in the case of uncorrelated boundaries the two rough surfaces are of the same statistical nature (see the factor 2 in the first two terms of Eq. (4.20b)).

The remarkable similarity between the behavior of the inverse attenuation length for both configurations can be observed in Fig. 2. Note that the normalization factor Λ_n for one boundary is twice as much of the quantity for two boundaries, and both curves are practically superposed.



FIG. 2: Plot of $\Lambda_n/2L_n$ versus kR_c for $kd/\pi = 100.5$, $n = 50$ and different values of $(k\sigma)^2$. Different sets of curves are shown for three waveguide configurations, as well as for the waveguide with one rough boundary analyzed in Ref. [17] (dashed line). Curves without symbols depict Eq. (6.4) for the waveguide with uncorrelated boundaries; curves labelled by triangles and diamonds stand for Eqs. (6.37) and (6.44) for waveguides with antisymmetric and symmetric boundaries, respectively

B. Waveguide with antisymmetric rough boundaries

According to Eqs. (4.28b) and (5.2), the inverse attenuation length can be presented as a sum of the partial inverse lengths $1/L_{n,n'}$ that are responsible for the scattering from nth into n'th modes,

$$\frac{1}{L_n} = \sum_{n'=1}^{N_d} \frac{1}{L_{n,n'}}. \tag{6.33}$$

However, it is noteworthy that in this case the partial terms describing the intramode $1/L_{n,n}$ and intermode $1/L_{n,n'\neq n}$ scattering are rather different since they are associated with different scattering mechanisms. Indeed, the intermode scattering results from the RHP spectrum only, through the expression in Eq. (4.26b) for the kernel $Q_1(k_x, q_x)$. These partial lengths are formed by the gradient scattering mechanism, due to the term containing in the scattering potential the derivative $\xi'(x)$, see Eq. (4.23a). They read as

$$\begin{aligned}\frac{1}{L_{n,n'\neq n}} &= 4\sigma^2 \frac{(\pi n/d)^2}{k_n d}\frac{(\pi n'/d)^2}{k_{n'} d}\sin^4[\pi(n-n')/2] \\ &\quad\times[W(k_n+k_{n'})+W(k_n-k_{n'})]. \end{aligned}\tag{6.34}$$

From the above expression one can draw an interesting conclusion. As one can see, in the case of symmetric boundaries the transitions are forbidden between the modes for which the difference of the indexes $n - n'$ is an even number. There are transitions only between odd and even channels. This fact can be explained by an existence of an additional integral of motion, namely, the parity. Specifically, if one solves the stationary problem for the effective Hamiltonian, there are two sets of eigenstates of different parity. Correspondingly, there are two independent sets of eigenvalues, see Refs. [16, 18].

The intramode scattering, related to the RSGP spectrum through the kernel $Q_2(k_x, q_x)$ in the Eq. (4.27b), is related to the SGS mechanism via the function $\hat{\mathcal{V}}(x)$ in Eq. (4.24a). Therefore, we have

$$\frac{1}{L_{n,n}} = \frac{\sigma^4}{2}\frac{(\pi n/d)^4}{k_n^2}\left[T(2k_n)+T(0)\right]. \tag{6.35}$$

Note that in a single-mode waveguide with $N_d = 1$ the sum over n' in Eq. (6.33) contains only one term with $n' = n = 1$. In this case the backscattering length $L_{1,1}^{(b)}$ entering in Eq. (6.35) is in accordance with that obtained in Ref. [13].

Bearing in mind very different dependencies of the $W(k_x)$ and $T(k_x)$ power spectra on the roughness correlation length R_c, one can write the inverse attenuation length as the sum of two terms. One is related to $W(k_x)$,

$$\frac{1}{L_n^{(1)}} = \sum_{\substack{n'=1,\\ n'\neq n}}^{N_d} \frac{1}{L_{n,n'}}, \tag{6.36}$$

and the other is just the inverse of the intramode attenuation length determined by Eq. (6.35). Therefore, $1/L_n^{(2)} = 1/L_{n,n}$, and one can write,

$$\frac{1}{L_n} = \frac{1}{L_n^{(1)}} + \frac{1}{L_n^{(2)}}. \tag{6.37}$$

In Fig. 2 we depict the length defined by Eq. (6.37) as a function of kR_c for different values of $k\sigma$; the curve is labelled with triangles.

As above, it is convenient to consider separately three regions of the correlation length.

1. *Small-scale roughness*

We start with the standard case of a small-scale boundary perturbation specified by the conditions (6.12). In this case for any term in the summation over n' the argument of the correlators $W(k_x)$ and $T(k_x)$ is much less than the scale of their decrease R_c^{-1}, therefore, it can be taken as zero. Thus, for the attenuation length describing the intermode scattering one gets,

$$\frac{\Lambda_n}{2L_n^{(1)}} \approx 4(k\sigma)^2\frac{n}{kd/\pi}\frac{W(0)}{k} \times \sum_{n'=1}^{N_d}\frac{(\pi n'/d)^2}{k_{n'}d}\sin^4[\pi(n-n')/2] \quad (6.38a)$$

$$\approx (k\sigma)^2\frac{n}{kd/\pi}\frac{kW(0)}{2} \quad (6.38b)$$

The factor $\sin^4[\pi(n-n')/2]$ gives alternating values of one and zero for $n-n'=1,3,..$ and $n-n'=2,4,..$, respectively, therefore, for a large number of conducting modes $N_d \gg 1$ it can be replaced by 1/2. The estimate (6.38b) is obtained with the change of the summation by integration. For a further analysis one has to take into account the formula in Eq. (6.14) for $W(0)$.

The second term in Eq. (6.37) reads as

$$\frac{\Lambda_n}{2L_n^{(2)}} \approx \frac{\pi^2}{2}\frac{(k\sigma)^4}{k_n d}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}, \quad (6.39)$$

where the explicit form for $T(0)$ can be taken as in Eq. (6.17).

It is easy to realize that the relationship between lengths defined by Eqs. (6.38b) and (6.39) is of the same order as in the case of uncorrelated boundaries, see Eq. (6.20). In this way one obtains the expression (6.21) for the crossing point $(kR_c)_{cr}$ at which the curves of $\Lambda_n/2L_n^{(1)}$ and $\Lambda_n/2L_n^{(2)}$ intercept each other.

2. *Large-scale roughness: weak correlations*

This intermediate situation is specified by Eq. (6.22). In Eq. (6.36) we replace $\sin^4[\pi(n-n')/2]$ by 1/2, and change the sum by the integral since the correlator $W(k_x)$ is a smooth function of n' (see the discussion before Eq. (6.23)),

$$\frac{\Lambda_n}{2L_n^{(1)}} \approx \frac{(k\sigma)^2}{\pi}\frac{n}{kd/\pi}\int_{-k}^{k}dk_x\sqrt{k^2-k_x^2}\,\frac{W(k_n-k_x)}{k}. \quad (6.40)$$

Note that Eq. (6.40) coincides with Eq. (6.24b) for the case of two uncorrelated boundaries.

For the length $L_n^{(2)}$ in the large-scale-roughness region, $kR_c \gg 1$, one should use the general expression (6.35) because of arbitrary value of the parameter k_nR_c. Its normalized version reads

$$\frac{\Lambda_n}{2L_{n,n}} = \frac{\pi^2}{4}\frac{(k\sigma)^4}{k_n d}\frac{n^3}{kd/\pi}\frac{T(2k_n)+T(0)}{k^3}. \quad (6.41)$$

3. *Large-scale roughness: strong correlations*

In this limit case specified by Eq. (6.26), the distance between wave numbers k_n and $k_{n\pm1}$ is much larger than the variation scale R_c^{-1} of the correlator $W(k_x)$, see Eq. (6.28). This means that the probability of the intermode ($n' \neq n$) transitions is exponentially small and the total attenuation length is mainly formed by the incoherent intramode ($n'=n$) scattering,

$$\frac{1}{L_n} \approx \frac{1}{L_n^{(2)}}. \quad (6.42)$$

For strong correlations the correlator $T(k_x)$ is a sharp function of its argument k_x. Thus, the term with $T(2k_n)$ can be neglected in comparison with $T(0)$, and one gets,

$$\frac{\Lambda_n}{2L_n} \approx \frac{\pi^2}{4}\frac{(k\sigma)^4}{k_n d}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}. \quad (6.43)$$

In this region of kR_c the SGS mechanism is recovered, determining the total attenuation length. The second inequality in Eq. (6.27) is automatically satisfied for any value of kR_c

C. Waveguide with Symmetric Rough Boundaries

As above, for this configuration it is convenient to write down the inverse attenuation lengths as a sum of two terms,

$$\frac{1}{L_n} = \frac{1}{L_n^{(1)}} + \frac{1}{L_n^{(2)}}. \quad (6.44)$$

Here each term is associated with the corresponding term in Eq. (4.36b), resulting from different mechanisms of the surface scattering.

The first term in Eq. (6.44) is related to the RHP spectrum $W(k_x)$ through the expression for $Q_1(k_x,k_x')$. According to Eq. (4.34b), we have

$$\frac{1}{L_n^{(1)}} = 4\sigma^2\frac{(\pi n/d)^2}{k_n d}\sum_{n'=1}^{N_d}\frac{(\pi n'/d)^2}{k_{n'}d}\cos^4[\pi(n-n')/2] \times [W(k_n+k_{n'}) + W(k_n-k_{n'})]. \quad (6.45)$$

The term $1/L_{n,n}^{(1)}$ corresponding to $n'=n$, is related to the amplitude scattering while the terms $1/L_{n,n'\neq n}^{(1)}$ result from the gradient scattering. These two mechanisms of surface scattering are due to the corresponding terms in the expression for $\Xi_1(k_x,k_x')$, see Eq. (4.32). Namely, the former follows from the term depending on the amplitude of the roughness profile $\sigma\xi_i(x)$, and the latter from the terms depending on the roughness gradient $\sigma\xi_i'(x)$.

The second term in Eq. (6.44) related to the RSGP spectrum through $Q_2(k_x,k_x')$, is associated solely with the SGS mechanism due to the square-gradient function $\widehat{\mathcal{V}}(x)$, see Eq. (4.33). In accordance with Eq. (4.35b), it is described by

$$\frac{1}{L_n^{(2)}} = \sum_{n'=1}^{N_d}\frac{1}{L_{n,n'}^{(2)}}. \quad (6.46)$$

Here

$$\frac{1}{L^{(2)}_{n,n}} = \frac{\sigma^4}{2}\frac{(\pi n/d)^4}{k_n^2}\left[\frac{1}{3}+\frac{1}{(\pi n)^2}\right]^2\left[T(2k_n)+T(0)\right] \tag{6.47}$$

and

$$\begin{aligned}\frac{1}{L^{(2)}_{n,n'\neq n}} =& \frac{128\sigma^4}{\pi^4}\frac{(\pi n/d)^2}{k_n}\frac{(\pi n'/d)^2}{k_{n'}}\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4}\\ &\times\cos^4[\pi(n-n')/2]\\ &\times\left[T(k_n+k_{n'})+T(k_n-k_{n'})\right].\end{aligned} \tag{6.48}$$

Note that in a single-mode waveguide with $N_d = 1$ (therefore with $n' = n = 1$ in Eqs. (6.45) and (6.46)) the backscattering length $L^{(b)}_{11}$ takes the expression derived in Ref. [15].

The diagonal part of the inverse attenuation length, $1/L_{n,n} = 1/L^{(1)}_{n,n} + 1/L^{(2)}_{n,n}$, characterizes the intramode scattering. The off-diagonal partial attenuation length $L_{n\neq n'} = 1/L^{(1)}_{n,n'\neq n} + 1/L^{(2)}_{n,n'\neq n}$ describes the intermode scattering.

To compare with the above case of the antisymmetric boundaries, here there are no transitions between the modes with odd values for the difference $n - n'$. Therefore, only the transitions between odd or even modes are allowed only. This fact is related to the underlying symmetry, and can be associated with specific integrals of motion of the total Hamiltonian.

1. *Small-scale roughness*

In the case of a small-scale roughness the first attenuation length reads

$$\frac{\Lambda_n}{2L^{(1)}_n} \approx 2(k\sigma)^2\frac{n}{kd/\pi}\frac{W(0)}{k}\sum_{n'=1}^{N_d}\frac{(\pi n'/d)^2}{k_{n'}d} \tag{6.49a}$$

$$\approx (k\sigma)^2\frac{n}{kd/\pi}\frac{kW(0)}{2}. \tag{6.49b}$$

For the second length we have,

$$\begin{aligned}\frac{\Lambda_n}{2L^{(2)}_n} \approx& \frac{\pi^2}{2}\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}\\ &\times\left\{\left[\frac{1}{3}+\frac{1}{(\pi n)^2}\right]^2+\frac{256k_nd}{\pi^4(\pi n/d)^2}\right.\\ &\times\left(\sum_{n'=1}^{n-1}+\sum_{n'=n+1}^{N_d}\right)\frac{(\pi n'/d)^2}{k_{n'}d}\frac{(n^2+n'^2)^2}{(n^2-n'^2)^4}\\ &\left.\times\cos^4[\pi(n-n')/2]\right\}\end{aligned} \tag{6.50a}$$

$$\approx \frac{\pi^2}{8}\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}. \tag{6.50b}$$

As well as in the configuration with uncorrelated boundaries, due to the estimate (6.16) every term in the sum of Eq. (6.50a) rapidly decreases with an increase of the absolute value of $\Delta n = n - n'$. In addition, the factor $\cos^4\pi(n-n')/2$ is zero for $n-n' = 1, 3, ...$ Therefore, the sum in Eq. (6.50a) can be evaluated by three terms with $n' = n, n\pm 2$, and one can replace the quantity in the curly braces by $1/4$. Here we also assume, $N_d \gg n \gg 1$.

It is evident that in the small-roughness case the crossing point is determined by Eq. (6.21).

2. *Large-scale roughness: weak correlations*

In the intermediate situation (6.22) the first attenuation length $L^{(1)}_n$ coincides with that obtained for the previous configurations,

$$\frac{\Lambda_n}{2L^{(1)}_n} \approx \frac{(k\sigma)^2}{\pi}\frac{n}{kd/\pi}\int_{-k}^{k}dk_x\sqrt{k^2-k_x^2}\,\frac{W(k_n-k_x)}{k}. \tag{6.51}$$

As for the second length $L^{(2)}_n$, for $N_d \gg n \gg 1$ it can be written,

$$\frac{\Lambda_n}{2L^{(2)}_n} \approx \frac{\pi^2}{16}\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)+T(2k_n)}{k^3}. \tag{6.52}$$

3. *Large-scale roughness: strong correlations*

In this region of kR_c the first attenuation length reads,

$$\frac{\Lambda_n}{2L^{(1)}_n} \approx 2\frac{(k\sigma)^2}{k_nd}\frac{n^3}{(kd/\pi)^3}kW(0). \tag{6.53}$$

For the SGS length one gets,

$$\frac{\Lambda_n}{2L^{(2)}_n} \approx \frac{\pi^2}{4}\frac{(k\sigma)^4}{k_nd}\frac{n^3}{(kd/\pi)}\frac{T(0)}{k^3}\left[\frac{1}{3}+\frac{1}{(\pi n)^2}\right]^2. \tag{6.54}$$

The ratio of the first attenuation length to the second one can be obtained due to Eq. (6.31). One can see that the amplitude scattering length always prevails over the SGS length within the interval of strong correlations, $L_n^{(1)}/L_n^{(2)} \ll 1$. For this reason, the condition of applicability coincides with Eq. (6.32).

In Fig. 2 by diamonds we show the behavior of the total dimensionless inverse scattering length as a function of the correlation parameter kR_c.

VII. NUMERICAL ANALYSIS OF THE SURFACE SCATTERING POTENTIAL $\widehat{U}$

In this Section we present numerical data for periodic waveguides (billiards) with the potential $\widehat{U}(x,z)$. Our main attention is payed to the structure of eigenstates for the discussed types of boundaries. Although this information is not directly related to the scattering properties of open waveguides, it is instructive to reveal a role of the SGS-terms in the potential, by comparing the structure of eigenstates for different boundaries. The numerical method is described in details in Ref.[18]; below we discuss the most interesting results and relate them to the previous studies (see, Ref.[16] and references therein).

Our particular interest is in the "repulsion" effect discovered in Ref. [18] for quasi-1D billiards with rough surfaces. This phenomena can be observed for a subset of eigenfunctions with the smallest values of the transversal quantum number of the unperturbed billiard with perfectly flat boundaries. For these eigenstates it was found that the maximum of the eigenfunction intensity is strongly shifted towards from the rough boundary. This "repulsion" was shown to increase with a decrease of the roughness correlation length R_c. Based on our numerical data, one can make a conclusion that this effect of "repulsion" is directly related to the SGS-terms discussed above.

To start with, we show in Fig. 3 a typical eigenstate of this type for the billiard with one rough surface (upper panel). To compare with, the low panel presents the corresponding eigenstate for the same potential, however, without SGS-terms. As one can see, by removing these terms, the repulsion disappears.

It is also instructive to compare the structure of such eigenstates with those computed for the case of both symmetric and antisymmetric rough boundaries, see Fig. 4. In both cases the potentials have the SGS-terms. However, the repulsion phenomena is absent in the antisymmetric billiard. A close inspection reveals that the difference between these two cases is entirely due to the fact that in the antisymmetric billiard there is no square-gradient scattering between different channels, in contrast to the symmetric billiard. Thus one can make a conclusion that the main contribution to the repulsion is due to the intermode SGS-terms. This conclusion supports an observation found in Ref. [18] according to which the effect of the repulsion can be explained due to a strong localization in the channel space. This localization occurs due to a relatively strong interaction between different conducting channels (waveguide modes).



FIG. 3: (Color online) Upper panel: intensity of a typical eigenstate manifesting a "repulsion" from the rough surface. Low panel: the corresponding eigenstate for the potential without the SGS-terms.

VIII. CONCLUSION

We have developed a perturbative approach to the surface scattering in multimode quasi-1D waveguides with rough surfaces. The approach is based on the construction of an effective Hamiltonian that formally corresponds to the "bulk" scattering described by the potential of a very complicated structure. The detailed analysis of this potential has led to the conclusion that there



FIG. 4: (Color online) Intensity of an eigenfunction revealing strong repulsion from two symmetric rough boundaries (low panel). The same for the antisymmetric billiard for which there is no interaction between different channels (upper panel).

are specific square-gradient scattering (SGS) terms that were not taken into account in the previous studies. We show that in spite of their seeming smallness in comparison with the standard terms, they can give a major contribution to the value of the scattering length in the case of a sufficiently strong roughness characterized by a corresponding value of the derivative of boundary profiles.

In order to demonstrate a strong influence of this SGS-mechanism, we have applied the approach to the waveguides with the boundaries of three different types. Specifically, we considered the waveguides with two uncorrelated boundaries, and compared the results with those obtained for other two cases of symmetric and antisymmetric rough boundaries. In this way one can see the influence of various terms very clearly.

According to our analysis, all terms in the expression for the effective potential can be regrouped in three parts. The first one is due to the roughness mean amplitude σ. Another group is characterized by the derivative of profiles, reflecting the fact that the scattering depends on two characteristic parameters, namely, on the amplitude σ (or, the variance σ^2), and on the correlation length R_c inversely proportional to the mean derivative of the profiles. In the expression for the scattering potential the above two types of terms have linear dependence on the root-mean-square roughness height σ. There is one more group of terms that can not be reduced to the two previous ones, and that gives rise to the square dependence on the profile derivative σ/R_c. For this reason, in the previous studies it was assumed that these SGS-terms can be safely neglected.

However, we show that apart from the σ-dependence due to the amplitude and gradient terms, as well as the σ^2-dependence due to the SGS-terms, there is also dependence on the correlation parameter R_c. As was found, the R_c-dependence for the SGS-terms turns out to be very different from those corresponding to the amplitude and gradient terms. Our analysis reveals that there is a region of parameters where the SGS-contribution to the inverse scattering length is compared (or even larger) with the standard contributions due to the amplitude and gradient scattering.

In accordance with the general theory of surface scattering [1] the wave propagation through any n-th conducting channel is determined by its total (or outgoing) attenuation length L_n. The inverse attenuation length $1/L_n$ can be presented as the sum of partial inverse lengths, see Eqs. (6.4), (6.37) and (6.44). The first term $1/L_n^{(1)}$ is associated with both the amplitude and gradient scattering, it depends on the roughness-height power spectrum $W(k_x)$. To the contrary, the partial length $L_n^{(2)}$ being connected with the SGS-terms, depends on the power spectrum $T(k_x)$ that is principally different from $W(k_x)$. For the convenience of comparison between three different types of boundaries, in Eq. (6.4) we have separately written the term $1/L_n^{(2)\updownarrow}$ that appears in the specific case of two uncorrelated boundaries. This term also depends on $T(k_x)$.

Such a representation allows us to analyze the total dependence of the inverse attenuation length $1/L_n$ on two control parameters, the roughness hight σ and the correlation length R_c. Our analysis shows that one can speak about three characteristic regions of R_c, namely, the small-scale (6.12), large-scale weak-correlation (6.22) and large-scale strong-correlation (6.26) roughness.

As we noticed above, the $W(k_x)$- and $T(k_x)$-power spectra have very different dependencies on the roughness correlation length R_c. Specifically, the inverse value of the L_2-length decreases with an increase of the correlation parameter kR_c; this occurs in the whole interval of the roughness values. As for the inverse value of the L_1-length, it shows a different behavior in each of the three regions of roughness and strongly depends on the waveguide configuration.

In the region of a small-scale roughness (6.12) the inverse length $1/L_1$ increases with an increase of kR_c. Therefore, the curves corresponding to the terms $1/L_n^{(1)}$ and $1/L_n^{(2)}$ $(1/L_n^{(2)} + 1/L_n^{(2)\updownarrow}$ for the case of uncorrelated boundaries) intersect each other, see Fig. 2. If the crossing point $(kR_c)_{cr}$ falls into this region of small-scale roughness, it obeys the law given by Eq. (6.21). To the left from the crossing point $(kR_c)_{cr}$, the SGS-length L_2 prevails over the standard length, $L_n^{(2)} \ll L_n^{(1)}$. Contrary, to the right from the point $(kR_c)_{cr}$, the L_1-length gives the main contribution to the scattering process, $L_n^{(1)} \ll L_n^{(2)}$.

The detailed analysis of three types of boundaries shows that for the small-scale roughness region one can specify that the crossover (with an increase of kR_c) occurs due to the following interplays:

- uncorrelated boundaries: the two SGS-terms that depend on the $T(k_x)$-power spectrum, versus the amplitude and gradient terms depending on the $W(k_x)$-power spectrum;
- antisymmetric boundaries: the SGS-term versus the gradient term;
- symmetric boundaries: the SGS-term versus the amplitude and gradient terms.

Thus, we show that in the region of small-scale roughness and for any fixed value of the roughness height σ, one can indicate the region of small values of the correlation length R_c where the SGS-length $L_n^{(2)}$ predominates over the the length $L_n^{(1)}$. This predominance occurs in spite of the fact that $1/L_n^{(1)}$ is proportional to σ^2 while $1/L_n^{(2)}$ is proportional to σ^4.

It is interesting to note that in the region to the left from the point $(kR_c)_{cr}$ we have observed that the length L_n^{SB} for symmetric boundaries is twice as much as the length L_n^{UB} for the uncorrelated boundaries. On the other hand, the latter length is twice as much as the length L_n^{AB} for the antisymmetric boundaries, i.e., $L_n^{SB} \approx 2L_n^{UB} \approx 4L_n^{AB}$.

There is another border that is characterized by the transition to the large-scale roughness with strong correlations, where a very different behavior of the total inverse length $1/L_n$ was found for the antisymmetric boundaries, in comparison with the uncorrelated and symmetric ones. For the two last cases, the major mechanism of the scattering is due to the amplitude terms in the potential, and a similar behavior with respect to kR_c $(1/L_n \propto kR_c)$ occurs for both of them. However, the attenuation length for the uncorrelated boundaries turns out to be twice as much as the length for the symmetric boundaries, i.e., $L_n^{UB} \approx 2L_n^{SB}$ (note that this relation is inverted in comparison with the region of small-scale roughness).

As for the waveguide with the antisymmetric boundaries for which the amplitude terms are absent in the potential, the SGS-mechanism prevails in this region, leading to the dependence $1/L_n \propto (kR_c)^{-3}$. Note, that here there is the interplay between the SGS-term and gradient one. The transition point was found to be $kR_c \sim k\Lambda_n$, with the crossover specified by the following interplays:

- uncorrelated and symmetric boundaries: the amplitude and gradient terms versus the amplitude terms,
- antisymmetric boundaries: the gradient and SGS terms versus SGS-term.

Finally, there is an intermediate region of a large-scale roughness and weak correlations where the first attenuation length mainly contributes, $L_n^{(1)} \ll L_n^{(2)}$. Here one can observe a strong similarity of the inverse attenuation length behavior for all configurations considered. It should be noted that this region is quite difficult for obtaining simple analytical estimates due to a strong dependence of the number of terms in the corresponding sums on the parameter kR_c. Numerical data manifest that the attenuation length roughly remains the same inside this intermediate region.

Acknowledgments

The authors acknowledge the support by the CONACYT (México) grant No 43730.

---

[1] F. G. Bass and I. M. Fuks, *Wave Scattering from Statistically Rough Surfaces* (Pergamon, New York, 1979).

[2] J. A. Konrady, J. Acoust. Soc. Am. **56**, 1687 (1974).

[3] A. R. McGurn and A. A. Maradudin, Phys. Rev. B **30**, 3136 (1984).

[4] Z. Tešanović, M. Jarić and S. Maekawa, Phys. Rev. Lett. **57**, 2760 (1986).

[5] N. Trivedi and N. W. Ashcroft, Phys. Rev. B **38**, 12298 (1988).

[6] A. B. Isers, A. A. Puzenko and I. M. Fuks, Akust. Zh. **36**, 454 (1990) [Sov. Phys. Acoust. **36**, 253 (1990)]; Journal of Electromagnetic Waves and Applications **5**, 1419 (1991).

[7] A. A. Krokhin, N. M. Makarov and V. A. Yampol'skii, J. Phys.: Condens. Matter **3**, 4621 (1991).

[8] V. I. Tatarskii, Waves in Random Media **3**, 127 (1993); *ibid* **7**, 557 (1997); *ibid* **10**, 339 (2000).

[9] A. E. Meyerovich and S. Stepaniants, Phys. Rev. Lett. **73**, 316 (1994); Phys. Rev. B **51**, 17116 (1995); Phys.

Rev. B **58**, 13242 (1998); Phys. Rev. B **60**, 9129 (1999); J. Phys.: Condens. Matter **12**, 5575 (2000).

[10] N. M. Makarov, A. V. Moroz and V. A. Yampol'skii, Phys. Rev. B **52**, 6087 (1995).

[11] A. M. Bratkovsky and S. N. Rashkeev, Phys. Rev. B **53**, 13074 (1996).

[12] G. A. Luna-Acosta, Kyungsun Na, L. E. Reichl and A. Krokhin, Phys. Rev. E **53**, 3271 (1996).

[13] N. M. Makarov and Yu. V. Tarasov, J.Phys.: Condens. Matter **10**, 1523 (1998).

[14] J. A. Sánchez-Gil, V. Freilikher, A. A. Maradudin and I. V. Yurkevich, Phys. Rev. B **59**, 5915 (1999).

[15] N. M. Makarov and Yu. V. Tarasov, Phys. Rev. B **64**, 235306 (2001).

[16] G. A. Luna-Acosta, J. A. Méndez-Bermúdez and F. M. Izrailev, Phys. Rev. E **64**, 036206 (2001).

[17] F. M. Izrailev, N. M. Makarov and M. Rendón, Phys. Rev. B **73**, 155421 (2006).

[18] J. A. Méndez-Bermúdez, G. A. Luna-Acosta and F. M. Izrailev, Physica E **22**, 881 (2004).

[19] F. M. Izrailev, N. M. Makarov and M. Rendón, Phys. Rev. B **72**, 041403(R) (2005); phys. stat. sol. (b) **242**, 1224 (2005).

[20] R. C. Bourret, Nuovo Cimento **26**, 1 (1962).